

COASTAL OCEAN RESEARCH

R/CZ-144: 3.1.1998-2.29.2000

Domoic Acid in Marine Diatoms: Biochemical Pathways and Environmental Regulation

G. Jason Smith

San Jose State University—Moss Landing Marine Laboratories

Background

he coastal United States has more toxic algal species, more algal toxins and more toxic algal blooms than ever before. The apparent spread of harmful agal blooms not only poses a human health threat but also has significant economic consequences for the seafood industry.

While declining water quality may be one reason for many of the observed increases in toxic blooms. the mechanisms through which man-made pollutants contribute to algal blooms is not well understood. Some of the "new" harmful algal blooms, for instance, have occurred in relatively pristine waters off the coast of Alaska and may be an artifact of better environmental testing. In the case of domoic acid, one of the more recently discovered toxins on the West Coast, sediment cores suggest the diatoms that synthesize the toxin have lived in coastal waters for more than a thousand years and are ubiquitous in coastal waters.

The Project

The purpose of this project was to identify environmental cues that can trigger toxic Pseudo-nitzschia blooms by examining the biochemical pathways of domoic acid (DA) production in these diatoms. Diatoms within the genus Pseudonitzschia produce DA.

More specifically, the focus of the study, led by Dr. G. Jason Smith of Moss Landing Marine Laboratory, was to investigate the hypothesis that DA is produced in Pseudonitzschia in response to stress. This hypothesis is motivated, in part, by the striking structural similarity between domoic acid and proline,

an amino acid produced by many marine organisms in response to stress.

The Method

To examine the effects of environmental stress on diatoms, Dr. Smith exposed diatoms to varying levels of copper, urea, nitrate and silicate. Copper is an essential nutrient that becomes toxic at high levels. Nitrate and urea both contain nitrogen, which is required for reproduction and cell growth. For this reason, plankton blooms cannot occur without a minimal amount of nitrogen. Silicate is another essential nutrient, used by diatoms to construct their frustules, or shells.

After exposing diatoms to nutrients, Dr. Smith measured levels of three key amino acids-DA, proline, and taurine. Taurine may reduce the toxicity of DA.

The Findings

The findings from this project support the theory that stress can trigger DA production. Dr. Jason Smith found that:



Photomicrograph of a dense bloom of diatoms from Monterey Bay, September 2000. The dominant needle-like cell chains are Pseudo-nitzschia australis. Photo: G. Jason Smith ©2000.



Photomicrograph of the toxic diatom Pseudonitzschia australis, isolated from a net tow during a bloom in Monterey Bay in September 2000. Note: the middle cell is alive while the two flanking cells are in states of senescence. Photo: G. Jason Smith ©2000.

 as silicate levels were lowered, DA production increased as much as a hundredfold:

 cells producing high levels of DA had relatively low levels of proline and relatively high levels of taurine; and.

 DA levels were higher when copper concentrations were higher and lower when copper levels were lower.

He also observed that nitrogen in urea and nitrate had no effect on DA production, beyond its known role in regulating cell growth.

Dr. Smith recently received funding from the National Science Foundation to isolate genetic markers that can be used to monitor DA production in diatoms from Monterey Bay.

Applications

This project helps answer basic genetic and biochemical questions about how and why cells produce DA. This information potentially can be used to help develop treatments for shellfish poisoning, and it has direct relevance for improving monitoring of coastal waters. At present, the state tests waters for the presence of Pseudo-nitzschia

diatoms but does not have the ability to tell whether these diatoms are actively producing DA. Dr. Smith said that his work lays the foundation for genetic fingerprinting that can discriminate between DAproducing and non-producing diatoms.

Cooperating Organizations

Merck & Company, Inc.

Monterey Bay Aquarium Research Institute

Provasoli Guilliard National Center for the Culture of Marine Phytoplankton

University of California, Santa Cruz—Marine Sciences Institute

Presentations

Ladizinsky, N.L., G.J. Smith, and K.H. Coale, 2000. Intracellular accumulation of domoic acid by *Pseudonitzschia* multiseries in response to increasing total copper levels. Sanctuary Currents 2000. Monterey Bay National Marine Sanctuary Symposium: Causes of Ecosystem Change: Natural or Human?, Santa Cruz, California.

Ladizinsky, N.L. and G.J. Smith, 2000. Accumulation of domoic acid by the coastal diatom *Pseudo-nitzschia* multiseries: A possible copper complexation strategy. PSA Annual Meeting, San Diego. Student Bold Award Presentation runner-up.

Smith, G.J. 1999. HAB or HAB NOT Workshop: Evaluation of the relationship between domoic acid and proline metabolism in *Pseudonitzschia* spp. from Monterey Bay, California. Invited instructor for an online, interactive presentation sponsored by the College of CoExploration and USC Sea Grant. Conference website: http:// www.coexploration.org/hab. Presentation website: http://members. aol.com/symbios222GJSHAB99.html.

Awards

The Fred and Ethel Meyers Marine Biology Trust Award Sigma Xi Grant in Aid of Research

Graduate Trainee and Thesis

Ladizinsky, Nicolas L., M.S. in Chemical Oceanography, California State University, Monterey Bay, anticipated June, 2002, "Accumulation of Domoic Acid by Diatoms in the genus *Pseudo-nitzschia* in Response to Increasing Cupric Ion Activity: A Possible Complexation Strategy."

For more information:

Dr. G. Jason Smith Environmental Biotechnology Lab Moss Landing Marine Laboratories San Jose State University Tel.: (831) 633.7270 Email: symbios@aol.com

PUB. NO. CSG-CZ-02-013

JANUARY 2002

California Sea Grant is a statewide, multiuniversity program of marine research, education, and outreach activities, administered by the University of California. Sea Grant-sponsored research contributes to the growing body of knowledge about our coastal and ocean resources and, consequently, to the solution of many marine-related problems facing our society. Through its Marine Extension Program, Sea Grant transfers information and technology developed in research efforts to a wide community of interested parties and actual users of marine information and technology, not only in California but throughout the nation. Sea Grant also supports a broad range of educational programs so that our coastal and ocean resources can be understood and used judiciously by this and future generations.

The national network of Sea Grant programs is a unique partnership of public and private sectors, combining research, education, and technology transfer for public service and dedicated to meeting the changing environmental and economic needs in our coastal, ocean, and Great Lakes regions.

This work is sponsored in part by a grant from the National Sea Grant College Program, National Oceanic and Atmospheric Administration, U.S. Department of Commerce, under grant number NA06RG0142, Project number A/P-1. The views expressed herein are those of the author and do not necessarily reflect the views of NOAA or any of its sub-agencies. The U.S. Government is authorized to reproduce and distribute for governmental purposes.



CALIFORNIA SEA GRANT COLLEGE PROGRAM Russell A. Moll, Director • Dolores M. Wesson, Deputy Director • Paul Olin, Interim Associate Director for Extension • Marsha Gear, Communications Coordinator University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0232 Phone:(858) 534-4440 Fax: (858) 453-2948 Web site: http://www-csgc.ucsd.edu